

Keladi Shivappa Nayaka University of Agricultural and Horticultural Sciences, Shivamogga, Karnataka

Gramin Krishi Mausam Sewa Project







Weather Forecast from 30.11.2022 to 04.12.2022

No. GKMSB/089/2022

Date: 29.11.2022 Weather Forecast Issued by the India Meteorological Department for Dakshina Kannada District for the coming five

days - until 0830 hrs of 30.11.2022 to 04.12.2022

Sl. No	Weather Parameters	Day-1 (30.11.2022)	Day-2 (01.12.2022)	Day-3 (02.12.2022)	Day-4 (03.12.2022)	Day-5 (04.12.2022)	Past Week Mean	Forecast Mean
1	Rainfall (mm)	11	0	0	3	2	48.2	16↓
2	Max Temp (°C)	29	30	31	31	31	31.4	30.4↓
3	Min Temp (°C)	24	24	24	24	24	22.4	24↑
4	Cloud cover (Okta)	6	5	5	6	6	4	6↑
5	Max. Relative Humidity (%)	88	86	86	88	88	89.8	87.2↓
6	Min. Relative Humidity (%)	75	73	73	75	75	65	74.2↑
7	Wind Speed (Kmph)	1	1	1	1	2	7	1.2↓
8	Wind Direction (°)	68	71	71	68	71	208	70 ↓
9	Forecast Warning	SCT	DRY	DRY	ISL	ISL		
10	Chance of Occurrence (%)	25-50	0	0	1-25	1-25		

As per the extended range rainfall forecast given by IMD, New Delhi and MC, Bangalore for coastal region of Karnataka including Dakshina Kannada district the rainfall will be Normal from 04.12.2022 to 10.12.2022

Summary of Weather Forecast for next Five days


As per the Weather forecast issued by the India Meteorological Department, Pune and Bengaluru for Dakshina Kannada district, Cloudy weather associated with a chance of Light to Moderate rainfall in most places of district, is likely to be expected for next five days. The maximum temperature would be around 29-31°C and minimum temperature would be around 24°C to during next five days. The Morning Relative humidity (RH) would vary from 86-88 per cent, Evening Relative humidity (RH) would vary from 73-75 per cent and wind speed will be in the range of 1-2 km/hr.

Mobile Applications developed by India Meteorological Department for the benefit of Farming Community Farmers are suggested to download the app using below links

	Mausam:	Meghdoot:	Damini:
Andriod:	https://play.google.com/store/app	https://play.google.com/store/apps/detail	https://play.google.com/store/apps/det
	s/details?id=com.imd.masuam	s?id=com.aas.meghdoot	ails?id=com.lightening.live.damini
Apple	https://apps.apple.com/us/app/id1	https://apps.apple.com/in/app/meghdoot/	https://apps.apple.com/app/id1502385
OS:	<u>522893967</u>	<u>id1474048155</u>	<u>645</u>

Decadal Rainfall (mm) Comparison:

=														
	Year	30 years Mean	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
	Rainfall	3790.4	3628.1	2984.2	3325.3	2816.3	2383.4	2827.1	3000	3381.2	4118.4	4110.8	4110.8	3519.5
	Painy days	110	128	105	127	112	107	110	121	112	123	126	152	117



Weather based Agromet Advisories for Dakshina Kannada District

Horticulture Crops:

Crop					
Crop	Operations/ Agromet Advisories				
General Advise	Farmers are advised to install Meghdoot, Mausam, Umang & Damini apps from mausam.imd.gov.in and be updated with weather events.				
Arecanut	To manage weeds in orchard spray Glyphosate (Amoniam salt)41%SL @ of				
(Weed	6ml/ltr or Glyphosate 54% SC @ 4ml/litre or Glyphosate 71% SC @ 6 gram per				
management)	litre.				
Arecanut	Farmers can Mulch the root zone of a crop with dried leaves or crop residues to				
(Water	save the moisture or to avoid the loss of water from root zone. Farmers should				
management)	irrigate the cropi.e., 20 litre per day per palm.				
Arecanut	Farmers can Spray Hexaconazole or Propiconazole or Difenconazole 1ml/ltr to				
(Leaf Spot)	the infected leaves.				
Coconut	Farmers can Mulch the root zone of a crop with dried leaves or crop residues to				
(Water	save the moisture or to avoid the loss of water from root zone. Farmers should				
management)	irrigate the cropi.e., 35 litre per day per palm.				
Coconut	To manage weeds in orchard spray Glyphosate (Amoniam salt)41%SL @ of				
(Weed	6ml/ltr or Glyphosate 54% SC @ 4ml/litre or Glyphosate 71% SC @ 6 gram				
management)	per litre.				
Banana (General Advice)	 5 months After planting spray Banana special @5gm/ltr of water at an interval of 30days or one month.(For one sprayer spray solution add 1 lime and shampoo) Bunch Treatment: To get good yield & quality bunches treat the bunch with 10 gram of Urea,10 gm of Potassium Sulphate & 0.5kg of cowdung should be mixed with 100ml of water & tie to end of bunch 				
Black Pepper	Farmers can Mulch the root zone of a crop with dried leaves or crop residues to				
(Water	save the moisture or to avoid the loss of water from root zone. Farmers should				
management)	irrigate the crop upto 20litre per day per vien.				
Jasmine	Right time to prune Jasmine plant. Diseased plant parts & dried shoots should be				
(Pruning)	pruned first. Pruned parts should be smeared with 10% Bordo paste				
Jasmine	For the Management of leaf spot disease, spray with Hexaconazole @ 1ml/litre				
(Leaf Spot)	of water.				
Cashew	Clean the infected place. After cleaning fill the infected part with				
(stem borer)	Chloropyriphos granules mixed with sand(1:1 ratio)				
Cashew	Farmers should spray Monocrotophos @ of 1.7ml/ltr or Lamdacyalothrin @				
(Tea Mosquito	0.5ml/ltr.				
Bug)	O.D.IIII/ICI.				

Cereals and Pulses:

Agronomic	After the harvest of paddy farmers can go for sowing of Dicot crops like Black
Advice	gram, Green gram, Beans and Green Manure seeds in the residual moisture.

Animal Husbandry:

Cattle	Animals should be tied at closed rooms or cattle sheds to protect them from cold.
(General advise)	➤ Cleanness should be maintained at Cattle shed & make sure of proper ventilation.
	➤ Right time for deworming.(contact nearest Veterinary officer for deworming
	tablets)
	> To avoid <i>Bowain Mastitits</i> Complete sucking of milk should be done without miss
	from Udder.

Mr. Praveen K.M , Technical Officer, GKMS, ZAHRS, Brahmavar, Udupi. Dr. K.V. Sudhir Kamath, Nodal Officer, GKMS, ZAHRS, Brahmavar, Udupi.